



Armed Forces College of Medicine AFCM





Nutritionally essential elements with their impacts on liver diseases



Dr. Marwa A. Dahpy



INTENDED LEARNING OBJECTIVES (ILO)



By the end of this lecture the student will be able to:

- Explain the biomedical importance of minerals
- Correlate copper metabolism to Wilson disease





**I-Macrominerals: potassium ,
Mg , sulfur**

**II-Trace elements: cu, selenium, zinc,
chromium,iodine,floride ,cobalt**

Copper metabolism

Biological functions:

Alterations of plasma copper:

**1-Cu Toxicity Wilson's disease
(Hepatolenticular degeneration)**

**2-Copper
Deficiency
(Menks
syndrome)**



Classification of minerals

**I-
Macrominerals:
potasssium , Mg
, sulfur**

What are the minerals?



- **Inorganic** elements
which are **essential**
nutrients.

- **Not** changed by
digestion or
metabolism.



Classification of minerals



Minerals are classified into 3 groups

1- Macrominerals (major elements)

- Present in great amounts in tissues.
- The daily requirements for each is >100 mg/day

EX: 7 elements

Ca , Mg ,
Phosphorous,
Na, K, Cl , Sulfur

2- Microminerals (trace elements)

- Present in minimal amounts in tissues
- The daily requirements for each is < 100 mg/day

EX: Iron ,Iodine ,
Copper, Cobalt , chromium, Mn

3-Extra trace elements <1
mg/day Molybdenenum ,
Selenium , Zn



1-Potassium

1. 2/3 of potassium is present in tissues and body fluids (potassium is **the main intracellular cations**)
2. 1/3 is present in the skeleton
3. **Regulation of osmotic and acid-base balance**
4. **Nerve and muscle excitability**
5. **Normalize heart rhythms**



Foods High in Potassium



Avocado



Banana



Potatoes



Spinach



Beans



Citrus juices



Fish



2-Magnesium



- **1-Proper nerve function & neurotransmission.**
- **2-Stabilize the APT co-substrate in enzymatic reactions.**
- **2-Regulate Muscle contraction**
- **3-Plays an important role in regulating the neuromuscular activity of the heart; maintains normal heart rhythm.**
- **Decrease magnesium level**
 - tremors, poor appetite, p
 - nystagmus and hypomagne



3-Sulfur



1. Enters in the formation of keratin of
2. Enter in **structure** Insulin.
3. The sulfur-containing amino acids
(**methionine, cystine, cysteine**)
4. sulfur-containing Vitamins: **biotin and thiamine**



Quiz



The following cause a decrease in serum phosphate EXCEPT::

- A. Vitamin D deficiency.
- B. Renal tubular disease.
- C. Hypoparathyroidism.
- D. Chronic alcoholism.
- E. Excessive use of antacids.

II-Trace elements



- <https://www.google.com/url?sa=i&source=images&cd=&cad=rja&uact=8&ved=0ahUf%3A%2F%2Fwww.alamy.com%2Fstock-photo-3d-graphics-essential-trace-elements-11v3Y&ust=1569031222500633&ictx=3&uact=3>





Copper metabolism

Biological functions:

Case scenario



A 15 years-old girl presented with **abdominal pain** and **diarrhea** for 3 days. She became **jaundiced** and provisional diagnosis of **hepatitis**.

#Lab. Findings :

- **Increased transaminases**
- **CBC anemia and thrombocytopenia**
- **The total serum copper level was decreased**
- **The free serum copper level was increased**
- **Serum Ceruloplasmin level < 20 mg/dl decreased**
- **Urine copper excretion/24h : increased**





What is your clinical impression?



**Wilson's disease :
genetic disease
due to accumulation of large
amount of copper**


**Genetic study revealed mutation of a gene
encoding a membrane bound copper transporting
ATPase**



Copper

Food Sources:

- Liver & kidney , Meats
- Legumes, leafy green vegetables.
- fruits
- Nuts
- Chocolate
- Cows milk is poor sou



Normal level;

(normal serum copper 10-22 $\mu\text{mol/L}$) = 100 - 200 $\mu\text{g/dl}$



Biological functions:

➤ **Help in iron absorption in Ferric state**

Hb synthesis (ALA synthase is a cu containing enzyme)

(anemia)

➤ **Bone formation □ synthesis of collagen**

➤ **Nervous tissue function**

➤ **component of many intracellular metalloenzymes**




Activity of many enzymes as:



1. **Superoxide Dismutase (SOD)**
2. **Lysayle oxidase** defective synthesis of collagen , defective bone and teeth , hypotonia, hyperelastist~~y~~ and osteoporosis
3. **Tyrosinase (Melan~~i~~ne synthesis)**
Hypopigmentation
4. **Cytochrome c oxidase** lack of ATP synthesis , lethargy and weakness
5. **Ferroxidase**
6. **Dopamine b-hydroxylase**
7. **Desaturase**




Dietary intake



Ceruloplasmin : α_2 globulin plasma pr , copper binding protein),







<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig1/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig2/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig3/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig4/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig5/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig6/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig7/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig8/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig9/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig10/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig11/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig12/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig13/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig14/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig15/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig16/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig17/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig18/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig19/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig20/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig21/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig22/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig23/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig24/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig25/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig26/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig27/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig28/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig29/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig30/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig31/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig32/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig33/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig34/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig35/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig36/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig37/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig38/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig39/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig40/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig41/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig42/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig43/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig44/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig45/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig46/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig47/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig48/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig49/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig50/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig51/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig52/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig53/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig54/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig55/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig56/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig57/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig58/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig59/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig60/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig61/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig62/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig63/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig64/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig65/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig66/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig67/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig68/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig69/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig70/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig71/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig72/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig73/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig74/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig75/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig76/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig77/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig78/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig79/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig80/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig81/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig82/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig83/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig84/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig85/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig86/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig87/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig88/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig89/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig90/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig91/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig92/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig93/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig94/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig95/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig96/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig97/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig98/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig99/>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710400/figure/fig100/>

Quiz




• Which is the primary route for elimination of copper from body?

a. Renal excretion

b. Liver sulphation

c. Biliary excretion

d. Sloughing of intestinal mucosal cell





Alterations of plasma copper

**1-Cu Toxicity
Wilson's
disease (Hepat
olenticular
degeneration)**

**2-Copper
Deficiency
(Menks
syndrome)**

Alterations of plasma copper:



1-Cu Toxicity

- **Dietry :rare**
- **Genetic : systemic Cu Overload**
Wilson's disease (hepatolenticular degeneration).
- **Increased serum free cu levels**
- **Decreased plasma ceruloplasmin**
- **This disease is characterized by accumulation of large amounts of copper in different tissues**



Wilson's disease (Hepatolenticular degeneration)



➤Cause:

It is a genetic disease (**autosomal recessive disorder**),

due to mutation of ATP7B gene

encoding membrane bound copper transporting ATPase

➤Biochemical basis of Wilson's:

- (1) Excessive copper absorption from the intestine.
- (2) Failure of Copper excretion via bile.
- (3) Failure to synthesize ceruloplasmin (Inadequate incorporation of copper into apoceruloplasmin)

➤Clinical picture:


It is characterized by accumulation of large amounts of copper in liver and different organs :

- Age of onset of symptoms range from 6 to 40 years

- Incidence 1:30000






Abnormal copper metabolism



https://www.google.com/imgres?imgurl=https%3A%2F%2Fimage.slides%3A&vet=10ahUKEwiU5-KPr97kAhURmhQKHZ_nAEHQMwhBKAawAA..il

olism&docid=fr06pdm0ns8B-M&tbnid=3kRUgqx21E7z_M





Hepatic presentations of Wilson Disease



More common in children than in adults


Vague symptoms & non specific Hepatitis and liver cirrhosis

Sever hepatic failure

Hepatic decompensation with ascites

Peripheral edema


Hepatic Encephalopathy



New Five Year Program





GIT Module





Neurological presentation

- Tends to occur in 2nd and 3rd decades or later
- defective formation of myelin sheath
- 3 main movement disorders
 - ❖ Dystonia
 - ❖ Tremors
 - ❖ parkinsonism



✖ lentiform nucleus of
the brain





Psychiatric disorders

- **20% of cases**
- **Loss of Emotional control**
- **Aggressive antisocial behavior**
- **Depression**






Ocular signs

Cornea green or golden pigments around the cornea

(Keyser- Fleisher ring)

Due to deposition of cu in basement membrane

Lense Sun flower cataract due to cu deposition in the lense






https://www.google.com/search?biw=1024&bih=489&tbm=isch&sa=1&ei=TjuEXZSIN5G0Up_Pg5gE&q=wilson%27s+disease&oeq=wils&gs_l=img..1.1.0i6712j0i8.1192640.1194208..1196470...0.0.0.420.1591.3-2j2.....0....1..gws-wiz-img.k-Yqw9XuWc8#





Laboratory diagnosis

- **Low Ceruloplasmin level** **< 10 mg/dl**
(N: 20-35 mg/dl)
- **High Urinary copper excretion rate** **>100mg/day**
(N: 20-50 ug/24hs).
- **High Free copper level** **> 25 ug/dl OR >3.9 umol/L**
(N: 8-12ug/dl = 1.3 :1.9 umol/L)
- **Low total copper level**
- **Liver biopsy : CU levels** **>250 ug/g of dry weight**
(N: 20-50 ug/g of dry weight)





Treatment:

Penicillinamine:

which:

- chelates copper
- and increase its excretion in urine.

Trientine induces urinary copper excretion

Zinc can block intestinal absorption of cu (only in pt without neurological manifestations).

Liver transplantation





2-Copper Deficiency

- **Dietary is rare**
- **Genetic :Menks syndrome**

(Menk's kinky or Steely hair disease)

- **It is a genetic fetal disease (X linked recessive disorder), due to mutation of ATP7A gene**
- **encoding for copper-transport protein which is a trans membrane protein**
- **This gene present in all tissues except liver (very little)**
- **important for regulating Cu transport from small intestine to the blood ,**
- **accumulation of Cu in intestine and kidney**
- **Poor distribution of Cu in other body cells.**



- **Menkes syndrome symptoms :**



- Brittle, kinky, steely, sparse, or tangled hair.
- Pudgy, rosy cheeks, sagging facial skin.
- Hypopigmentation.
- Feeding difficulties.
- Irritability.
- Lack of muscle tone, floppiness.
- Low body temperature.
- Intellectual disability and developmental delay.
- Fetal disease



<https://www.google.com/url?sa=i&source=images&cd=&cad=rja&uact=8&ved=0ahUKEwjohP3Wr6vmAhUC3BoKHdy1C4YQMwhAKAAwAA&url=https%3A%2F%2Fghr.nlm.nih.gov%2Fcondition%2Fmenkes-syndrome&psig=AOvVaw0Y7LaGSju22IHov0W-POyA&ust=1576077319007358&ictx=3&uact=3>

- **OTHER COPPER DISEASES**

- 1. Idiopathic Copper Toxicosis
- 2. Tyrolian Infantile Cirrhosis
- 3. Indian Childhood Cirrhosis



Quiz



- Which is the treatment of Menkes disease ?
- ☐ A. Zinc
- ☐ B. Penicillamine
- ☐ C. Trientine hydrochloride
- ☐ D. Early parenteral copper administration and supportive measures



II-Other Trace elements: selenium, zinc,

chromium,iodine,floride ,cobalt

2-Selenium



1- is an important antioxidant .

2- It is important for many enzymes as

Glutathione Peroxidase.

3-Present in 25% of human proteins

Selenocysteine amino acid



3-Zinc



1- It is a cofactor of some enzymes as **Alkaline phosphatase & superoxide dismutase.**

2-Zn Finger motifs binds to DNA regulating gene expression

3-It is important for **storage of insulin**

4-wound healing & healthy skin.

5-Strong immune system

6-Antioxidant nutrient.

7-Healthy hair




4-Fluoride



Present in Drinking water & seafood

is present in the body almost entirely in bone and teeth.





5-Chromium

- It is an important as a component of **glucose tolerance factor (GTF)** which is a **factor** **potentiates** the effect of insulin by facilitating its binding to **the cell receptor**.
- **Reduce** cholesterol & t



6-Iodine



- **Present in:** Sea water ,iodized table salt & Sea foods
- **Enter in the formation of thyroid hormones**
- **Deficiency :excess stimulation of thyroid gland by TSH (Goiter)**

•7-Cobalt

- **Component of Vit B₁₂ (Methylcobalamin or Adenosylcobalamin)**

•8-Molybdenum

- **Essential component of some flavoprotein enzymes as xanthine oxidase and sulfite oxidase which catalyzes the oxidation of sulfite (SO₃⁻) to sulfate (SO₄⁻)**



Minerals act as antioxidants




- **Copper**
- **Manganese**
- **Zinc**
- **Iron**
- **Selenium**



Quiz



• Which of the following mineral deficiency may results into impaired growth and development, skin lesions and loss of appetite



- 1- Zinc
- 2- Cobalt
- 3- Magnesium
- 4-chromium



-Selenium is important for the following enzyme:



- A) Superoxide dismutase
- ☒ B) Alkaline phosphatase
- C) Glutathione peroxidase
- D) catalase





Lecture Summery

- **Inorganic mineral elements** that have a function in the body must be provided in the diet. When intake is insufficient, deficiency may develop.
- **Wilson's disease (hepatolenticular degeneration).**
Increased serum free cu levels ,Decreased plasma ceruloplasmin,
This disease is characterized by accumulation of large amounts of copper in different tissues
- **Genetic :Menks syndrome**
 - Accumulation of Cu in intestine and kidney ,Poor distribution of Cu in other body cells.

SUGGESTED TEXTBOOKS



1. Lippincott's Illustrated Reviews in Biochemistry





***Thank you
Marwa Dahpy***

